


CONTRACEPTIVE IMPLANT INFORMATION SHEET



• Keep the smaller bandage on for 3-5 days.

	When To Consult A Health Care Provider
	If you see any part of the implant protruding from the wound
	If you notice redness, warmth, drainage, persistent pain at the insertion site
	and have a fever (38°C)
	If any point you are NOT able to feel the implant
	You can call us at 604-736-7878 or go to your regular health care provider.



CONTRACEPTIVE IMPLANT INFORMATION SHEET

When Does It Start Working?

Immediately post-pregnancy – The implant is effect right away

Any other time – The implant takes 7 days to become fully effective it is advised that a secondary birth control method (ie, condoms) is used during this time.

What Will Happen To My Period?

You may experience longer/shorter periods or have no bleeding at all. The time between your periods may change and you might experience spotting.

How It is Removed?

It is your responsibility to have the implant removed or replaced in 3 years time.

The removal process is quick and easy, your skin is numbed and a small incision is made and the implant is removed at any office or clinic at anytime.

Is It Easy To Get Pregnant After?

Most people get pregnant easily after the implant is removed.

Possible Side Effects:

- □ Mood changes
- U Weight gain
- Headaches
- □ Acne, and/or skin
 - changes in the upper arm
- □ Irregular bleeding

Side effects may go when the implant is removed.



Don't Forget Condoms!

Implants do not protect you from sexually transmitted infections.

